



## Ecology

# Effectiveness of two sampling methods for social wasps in different ecosystems

Mateus Aparecido Clemente<sup>1</sup>, Roger Guevara<sup>2</sup>, Hugo Ribeiro Moleiro<sup>1</sup>, Orlando Tobias Silveira<sup>3</sup>, Marcos Magalhães de Souza<sup>4</sup> & Edilberto Giannotti<sup>1</sup>

1. Universidade Estadual Paulista, UNESP, Rio Claro, São Paulo, Brazil. 2. Instituto de Ecologia do México, INECOL, Xalapa, Veracruz, Mexico. 3. Museu Paraense Emílio Goeldi Belém, Pará, Brazil. 4. Instituto Federal de Educação, Ciência e Tecnologia do Sul de Minas, Inconfidentes, Minas Gerais, Brazil.

EntomoBrasilis 14: e918 (2021)

### Edited by:

William Costa Rodrigues

### Article History:

Received: 09.vii.2020

Accepted: 07.xii.2021

Published: 20.i.2021

### Corresponding author:

Mateus Aparecido Clemente

✉ mateus1981@gmail.com

### Funding agencies:

↳ Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES); Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)

**Abstract.** There are different methodologies used to make an inventory of social wasps. In general, these methods are divided into active search and passive collections. Each method has a different performance, depending on the environment in which the collection is being carried out. Thus, the choice for the proper methodology according to the study area will impact the success of sampled species. This study aimed to evaluate the performance of sampling methodologies for social wasps (active search and bait traps) in different phytophysiognomies (Regenerating Cerrado, "Cerradão", Restored Forest, Semideciduous Forest and Riparian Forest) in the state of São Paulo, Southeastern Brazil. The active search sampled a greater number of species, with the exception of the Regenerating Cerrado. Regarding abundance, the active search was also more efficient, except in "Cerradão", where there was no difference between the methods, and in Regenerating Cerrado, where the traps sampled a higher number of individuals. The fact that none of the methods collected all the sampled species indicates that it is appropriate to use more than one collection method in order to obtain a sampling closer to the true richness of the studied sites.

**Keywords:** Active search; attractive traps; diversity; phytophysiognomies; richness.

In zoological inventories, ecology or taxonomy of social wasps, insects popularly known in Brazil as "marimbondos" or "cabas", different active and passive sampling methods are used for collection (SOUZA & PREZOTO 2006; SOUZA *et al.* 2015; SOMAVILLA *et al.* 2019).

The active methods are dependent of the collector presence in the field. There are different active sampling methods, such as: (1) Punctual sampling, when the collection is made in specific areas, such as floral resources (MARQUES *et al.* 2005; HERMES & KÖHLER 2006; CLEMENTE *et al.* 2013); (2) Utilization of quadrants, in which an area is pre-determined before the collection and then carefully examined (SOUZA & PREZOTO 2006); (3) Active search, when the collection is made by moving through an area in search for individuals or colonies, with the aid of an entomological net, with this methodology being the most used in social wasps samplings (DINIZ & KITAYAMA 1994; SILVEIRA 2002; ELPINO-CAMPOS *et al.* 2007; SANTOS *et al.* 2007; RIBEIRO-JUNIOR 2008; JACQUES *et al.* 2012; SILVA 2012; SOMAVILLA *et al.* 2014; SOUZA *et al.* 2020); (4) Active collection using a liquid bait of sucrose solution, with the aid of a pulverizer which sprays the solution over marked points on the vegetation of a trail (modified from LIOW 2001; LIMA *et al.* 2010; TANAKA-JUNIOR & NOLL 2011; LOCHER *et al.* 2014).

In the passive methods, traps are positioned and remain in field and therefore they have no direct interference of the collector; traps are methods in which an equipment is made to cease the insect's movement, in a way that when an insect enters the trap, it can no longer escape (ALMEIDA *et al.* 1998). These traps can be either flight interceptors, such as Malaise trap (SILVEIRA 2002; MORATO *et al.* 2008; SOUZA *et al.* 2015,

SOMAVILLA *et al.* 2019) or bait traps (LOCHER *et al.* 2014). The bait used in the traps can be of animal source, (such as sardines) (SILVEIRA *et al.* 2005; RIBEIRO-JUNIOR 2008; CLEMENTE 2009; TOGNI 2009), sugary liquids (SANTOS 1996; RIBEIRO-JUNIOR 2008; SOUZA *et al.* 2015), sodas (WEGNER & JORDAN 2005), chemical attractors (LANDOLT *et al.* 2000) and light traps used to attract species of Apoica (NETO *et al.* 1995).

Despite the rising number of studies regarding sampling methods of social wasps, there are few studies dealing with the efficiency of different methods when they are used in areas which have different ecosystems (SOUZA & PREZOTO 2006). Considering that a species inventory of a given area is the first step for its conservation and the rational usage of its biotic resources (MELO *et al.* 2005), the definition of a sampling methodology is a strategic tool to obtain secure information of the biota, hence justifying studies with this theme.

Thus, the present study aims to assess the efficiency of different sampling methodologies for social wasps' collection (active search and bait traps) in different ecosystems: Regenerating Cerrado, "Cerradão", Restored Forest, Semideciduous Forest and Riparian Forest).

## MATERIAL AND METHODS

The collects were made in five ecosystems of different phytophysiognomies in the State of São Paulo (Figure 1).

(A) A fragment of Semideciduous Forest (PAGANO & LEITÃO-FILHO 1987), with approximately 230 ha, having parts of both Rio Claro and Araras municipalities (22°21'06" S, 47°29'07" W), located at 630 m of altitude. The area has a semi-dense

canopy, with open canopy areas resulted from different levels of environmental stress, such as near sugar-cane plantations, fires and deforestation. The tree component has two floristically distinct stratifications: one from seven to 15 m high and another reaching 20 to 25 m high (PAGANO *et al.* 1987).

(B) An area of Restored Forest, located at the municipality of Iracemápolis (22°34'34" S, 47°30'25" W), at 624 m high. The restored area has approximately 20 ha and had its restoration process initiated in 1987 (SIQUEIRA 2002). The area is surrounded by sugar-cane plantations and is possible to observe different levels of canopy stratification, with a mean of 10 m high.

(C) An area of Regenerating Cerrado (22°24'22" S, 47°32'18" W), at 650 m high. The area is a local public location, with the southern border being limited by the *Universidade Estadual Paulista* (UNESP) and the northeastern border limited by the *Floresta Estadual Edmundo Navarro de Andrade* (FEENA), a fragment of "Paludosa" forest (a swamp-like environment), and an urban neighborhood which causes intense pressure. This vegetation area has approximately 100 m<sup>2</sup>, with a grove-like aspect, absence of underbrush and trees with height relatively uniform. The litter is scarce or inexistent, as the same as presence of seedlings (CARDOSO-LEITE *et al.* 2004).

(D) Riparian forest area (FRAGOSO 2005) located at the municipality of Itirapina, along the Lapa stream (22°22'16" S, 47°47'16" W), at 656 m of altitude, with an area of approximately 14 km<sup>2</sup>. The study site, originally part of a larger fragment of Mesophile Semideciduous Forest, suffers from strong agricultural impact, which reduced its original area to a thin band of riparian forest surrounded by sugar-cane plantations.

(E) "Cerradão" (22°24'49" S, 47°45'32" W), located at 671 m high, at the municipality of Ipeúna, is a phytophysiognomy which have some similarities with a "Cerrado" in a restricted sense. This area has a continuous canopy and arboreal coverage between 50% to 90%, directly influencing in luminosity conditions, thus favoring differentiations in the bushy and herbaceous strata. However, in the surrounding areas there are extensive sugar-cane plantations with burning practices during the pre-harvest, generating strong negative pressure towards the fragment. Additionally, a strong motorcycle traffic also occurs withing the fragment, causing damages to the ground and seedlings. Finally, several aggradations and tree falls were observed.

The field work was done from October 2011 to June 2013. During the whole year, in alternating months, two field trips were conducted in each area. The first trip of each month aimed to mount and set the traps and the second to disassemble and remove them. The first and second trip have a seven-days interval. Thus, each trap remained for a total of 1,008 h in the field, totalizing 10,080 h of passive collection per area.

Additionally, active searches were conducted during the same days of trap mounting/disassembling. The searching was compounded by two members which remained in field during seven hours/day, from 09 a.m. to 16 p.m., a time period with a higher foraging activity range of some groups of social wasps in different ecosystems (SANTOS & PRESLEY 2010; BARBOSA *et al.* 2014; BRITO *et al.* 2020). Thus, a total of 168 h of sampling effort per area was done in a year. The searching for individuals were conducted in local trails of each area, with the verification of nest occurrence in tree holes, large-leaved plants and edifications. The collected wasps were killed in recipient containing ether and were posteriorly stored in 70% alcohol.

The bait-trap distribution was done by the following procedure: ten points, distanced 100 m from each other, were marked using plastic tape tied to trees, thus reducing the chance of pseudoreplication by collection of a single wasp population in different sampling unities. At each point, five traps with a height of 1.5 m from ground and five other traps at canopy level (around 5-9 m) were set. The traps were made with 2 L plastic bottles (Souza *et al.* 2015). At each bottle, four circular holes were done and 200 ml of concentrated passion fruit juice were added to be used as bait. The specimens were collected with a sieve and tweezers and fixed in 70% alcohol recipients of the universal collector type.



**Figure 1.** Distribution map of the collection areas in the São Paulo State and the sampled spots in each phytophysiognomy. The yellow marking represents the distance between sampling points (100 m) and where each trap was set.

The sorting and pinning of specimens were done at the Rio Claro Campus of the UNESP. The species were identified by comparison with specimens deposited at the social wasp collection of the Zoology Department and by using identification keys to genera and species (RICHARDS 1978; COOPER 1997; CARPENTER & MARQUES 2001; PICKETT & WENZEL 2007).


The software used in the analyses were PAST - v. 1.49 (HAMMER *et al.* 2001), BioEstat v. 5.0 (AYRES *et al.* 2007) and R (R DEVELOPMENT CORE TEAM 2009).

The relative abundance of each species by each different method was measured by the ratio abundance of each species/total abundance. To verify which method was more efficient in each of the five sampled areas, the richness and abundance values were submitted to a linear mixed model test (lmm), aiming to verify the presence of significant differences. The type of vegetation, with five levels, and the sampling methods were set as fixed parameters. As each sampling point were repeatedly sampled, despite the method was changed, we used the identity of the points as a random parameter. It allowed us to infer the effects of microenvironments over the data variation in each kind of vegetation. For specific comparisons between sampling methods, we used the contrast based on the linear predictor of the model. To assess if the number of collections was enough to each sampling method (active search, bait traps) the estimators Jackknife 1 and 2 were applied.

## RESULTS AND DISCUSSION


The mixed model revealed significant differences at the

species richness catches by each method among all the vegetation kinds ( $\chi^2 = 19.4$ , g.l. = 4,  $p < 0.001$ ). In general, we determined that the active search has sampled a higher species richness, with exception of the Regenerating Cerrado, where the method resulted in 70% less species richness (mean = 3.5 species) in comparison to traps (mean = 6.3 species) at each sampling point  $z = 2.8$ ,  $p = 0.014$  (Figure 2).



**Figure 2.** Mean values of the sampled species richness per point in each phytophysiognomy in relation to the sampling method (active search and bait traps). Studied areas (RC - Regenerating Cerrado, C - Cerradão, RTF - Restored Forest, SDF - Semideciduous Forest and RPF - Riparian Forest).

In general, the abundance was more sampled by the active search as well, with higher absolute values of individuals (Figure 3). This difference was observed in three out of the five vegetation types ( $z > 3.1$ ,  $p < 0.004$ ): Restored Forest (172%), Semideciduous Forest (67%) and Riparian Forest (56%). On the other hand, there was no significant differences for the Cerradão, whilst for the Regenerating Cerrado we observed that traps resulted in a higher social wasp abundance (116%) in relation to the one resulted from active search ( $z = 6.2$ ,  $p < 0.001$ ).



**Figure 3.** Mean values of the sampled species abundance per point in each phytophysiognomy in relation to the sampling method (active search and bait traps). Studied areas (RC - Regenerating Cerrado, C - Cerradão, RTF - Restored Forest, SDF - Semideciduous Forest and RPF - Riparian Forest).

For the Regenerating Cerrado, differing from remaining areas, the most effective sampling method was bait trap, collecting 17 species (80.35%), in comparison to the active search (13 species - 61.9%) (Table 1). The same was verified for the abundance, resulting in 203 individuals captured (68.35%), whilst only 94 were captured by active search (31.65%). When compared to other areas of less anthropic pressure of this study, the Regenerating Cerrado resulted in

the highest abundance and the second highest social wasp richness among the sampled phytophysiognomies.

The Regenerating Cerrado has a degraded vegetation and a poor resource availability (CARDOSO-LEITE *et al.* 2004), hence turning the traps filled with passion fruit juice into attractive resources, what would explain the abundance values. Regarding species richness, the Regenerating Cerrado has in its surroundings a "Paludosa" forest and an area of eucalyptus planting in different successional stages (FEENA), which may house colonies of social wasps that may forage at the sampled area (RODRIGUES & MACHADO 1982).

At the FEENA, RODRIGUES & MACHADO (1982) made a twelve-year study and recorded 33 species and 10 genera of social wasps, demonstrating the expressive richness of this forest, close to studied site. Studies carried out in areas with a certain degree of impact, in Brazil, also resulted similarly (MARQUES *et al.* 1993; LIMA *et al.* 2000; MARQUES *et al.* 2005; ALVARENGA *et al.* 2010; AUAD *et al.* 2010; SANTOS & PRESLEY 2010; DE SOUZA *et al.* 2011; BUENO *et al.* 2019).

A total of 14 species were captured at the Cerradão, out of which 12 were captured actively (85.71%) in a total of 75 individuals (57.25%) (Table 1). This was the sole area in which bait traps and active search resulted in equal efficiency in relation to number of individuals collected, although it is concluded that for sampling richness, the active search was more effective. In relation to the phytophysiognomy, the Cerradão has a canopy predominantly continuous with coverage ranging from 50% to 90% (RIBEIRO & WALTER 1998), with canopy closure (RIZZINI 1997), fact which directly affects the active collects made inside the vegetation. Thus, it was expected that in the Cerradão the number of collected species by active search would be similar to that of the Semideciduous Forest, as it really happened in the present study (Table 2).

Other works show that transition areas between Cerrado and forest, as well as in more bushy physiognomies of Cerrado (such as Cerradão), usually have a higher richness of social wasps in relation to more open areas and adjacent agriculture areas. This may be explained by the larger structural complexity and availability of nesting sites (SANTOS *et al.* 2009; FERREIRA *et al.* 2020).

At the Restored Forest, 15 species were recorded, out of which 13 (86.67%) were collected by active search and nine (60%) by passive methods (Table 1). With an entomological net, 102 individuals (62.58%) were collected, whilst 61 (37.42%) were captured by the bait trap (Table 1).

No studies were found dealing with social wasps in Restored Forests, however, due to the hydric resources close to the sampling area and a quite heterogenous vegetation, it was observed that this environment is capable of maintaining relatively high populations of these animals.

At the Restored Forest, the methodology of active searching resulted in being more effective to both species richness and abundance (Table 2). It may be explained by the large spacing between trees and absence of canopy closure, as it kept the trails always illuminated thus facilitating the visual location of the wasps (RAVERET-RICHTER 2000).

At the Semideciduous Forest, 12 social wasp species were recorded, with 10 being collected by active search and eight (66.67%) by bait traps. In relation to abundance, 56 (73.68%) individuals were collected actively and 20 (26.32%) collected in traps (Table 1). This is the phytophysiognomy with the shortest proportion of collected species in relation to the estimation made to this ecosystem (Table 2). In

**Table 1.** Absolute abundance (Ab) and relative (F%) of wasp species sampled by active search (AS) and bait traps (BT) in the five studied areas.

Sampled physiognomies Abundance (Ab) and Frequency (F)	RC						C						RTF						SDF						RPF						Total
	BT		AS		BT		AS		BT		AS		BT		AS		BT		AS		BT		AS		BT		AS		BT		
	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	Ab	F%	
<i>Agelata multipicta</i>	10	4.93	0	-	1	1.79	3	4.00	1	1.64	2	1.96	5	25.00	4	7.14	2	2.08	6	4.00	34										
<i>Agelata pallipes</i>	117	57.64	61	64.89	14	25.00	17	22.67	6	9.84	7	6.86	1	5.00	18	32.14	39	40.63	18	12.00	298										
<i>Agelata vicina</i>	16	7.88	0	-	20	35.71	24	32.00	3	4.92	3	2.94	9	45.00	23	41.07	38	39.58	74	49.33	210										
<i>Apoica gelida</i>	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1	1.79	0	-	0	-	1										
<i>Apoica pallens</i>	1	0.49	1	1.06	1	1.79	0	-	3	4.92	0	-	0	-	0	-	2	2.08	1	0.67	9										
<i>Brachygastra lecheguana</i>	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1	1.04	3	2.00	4										
<i>Mischocyttarus drewseni</i>	0	-	1	1.06	0	-	0	-	0	-	4	3.92	0	-	1	1.79	0	-	15	10.00	21										
<i>Mischocyttarus mattogrossoensis</i>	2	0.99	8	8.51	0	-	0	-	0	-	2	1.96	0	-	2	3.57	0	-	1	0.67	15										
<i>Mischocyttarus rotundicollis</i>	0	-	1	1.06	7	12.50	14	18.67	0	-	21	20.59	0	-	0	-	0	-	2	1.33	45										
<i>Mischocyttarus paraguayensis</i>	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1	0.67	1										
<i>Mischocyttarus ignotus</i>	0	-	0	-	0	-	0	-	0	-	3	2.94	0	-	0	-	0	-	1	0.67	4										
<i>Mischocyttarus tricolor</i>	0	-	2	2.13	0	-	0	-	0	-	3	2.94	0	-	0	-	0	-	0	-	5										
<i>Mischocyttarus montei</i>	1	0.49	1	1.06	0	-	0	-	4	6.56	50	49.02	1	5.00	0	-	1	1.04	0	-	58										
<i>Polistes cinerascens</i>	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	3	2.00	3										
<i>Polistes billiardieri</i>	0	-	0	-	1	1.79	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1										
<i>Polistes lanio</i>	4	1.97	3	3.19	0	-	0	-	0	-	3	2.94	0	-	2	3.57	0	-	0	-	12										
<i>Polistes similimus</i>	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1	0.67	1										
<i>Polistes geminatus</i>	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1	0.67	1										
<i>Polistes subsericus</i>	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1	1.04	1	0.67	2										
<i>Polistes versicolor</i>	0	-	0	-	0	-	2	2.67	0	-	0	-	0	-	0	-	1	1.04	0	-	3										
<i>Polybia chrysothorax</i>	7	3.45	3	3.19	0	-	1	1.33	39	63.93	2	1.96	0	-	0	-	1	1.04	1	0.67	54										
<i>Polybia dimidiata</i>	7	3.45	4	4.26	2	3.57	1	1.33	1	1.64	0	-	1	5.00	1	1.79	4	4.17	1	0.67	22										
<i>Polybia fastidiosuscula</i>	3	1.48	0	-	8	14.29	8	10.67	2	3.28	1	0.98	1	5.00	3	5.36	1	1.04	10	6.67	37										
<i>Polybia ignobilis</i>	17	8.37	7	7.45	0	-	1	1.33	0	-	0	-	0	-	0	-	2	2.08	3	2.00	30										
<i>Polybia minarum</i>	0	-	1	1.06	0	-	0	-	0	-	0	-	1	5.00	1	1.79	0	-	0	-	3										
<i>Polybia occidentalis</i>	3	1.48	0	-	2	3.57	1	1.33	0	-	0	-	0	-	0	-	2	2.08	4	2.67	12										
<i>Polybia paulista</i>	5	2.46	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1	1.04	1	0.67	7										
<i>Polybia sericea</i>	3	1.48	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	3										
<i>Polybia jurinei</i>	5	2.46	0	-	0	-	1	1.33	2	3.28	1	0.98	1	5.00	0	-	0	-	0	-	10										
<i>Protonectarina sylveirae</i>	1	0.49	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1										
<i>Synoecca cyanea</i>	1	0.49	1	1.06	0	-	2	2.67	0	-	0	-	0	-	0	-	0	-	2	1.33	6										
<b>Total</b>	<b>203</b>	<b>100</b>	<b>94</b>	<b>100.00</b>	<b>56</b>	<b>100.00</b>	<b>75</b>	<b>100.00</b>	<b>61</b>	<b>100.00</b>	<b>102</b>	<b>100.00</b>	<b>20</b>	<b>100.00</b>	<b>56</b>	<b>100.00</b>	<b>96</b>	<b>100.00</b>	<b>150</b>	<b>100.00</b>	<b>913</b>										

general, a higher success during active search was obtained when the collectors got close to borders or opened areas, with higher luminosity, facilitating the visual location of the insects, as in RAVERET-RIECHTER (2000). The height of the trees and a too-closed canopy provide a low level of luminosity, hindering the collection with entomological net inside the phytophysiognomy, as in other related studies (DINIZ & KITAYAMA 1998; LIMA 2008; TANAKA-JUNIOR & NOLL 2011; SOUZA *et al.* 2012; TOGNI *et al.* 2014). Regarding the low efficiency of bait traps in this ecosystem, it is probably explained by its considerable heterogeneity, providing abundant nectar resources, turning the baits into less attractive resources.

The work of KLEIN *et al.* (2015), carried out in deciduous forests of the State of Rio Grande do Sul, also reported a low species richness and abundance collected by bait traps when compared to more open environments, which are scarcer in resources, such as monoculture borders close to forests. Therefore, it is suggested that for these environments, other sampling methods should be used, such as Malaise trap (SOMAVILLA *et al.* 2019).

The highest richness (23 species) was obtained at the Riparian Forest, with 21 (91.3%) collected by active search and 14 (60.87%) by bait traps. The same was observed for the abundance, in which 150 (60.98%) specimens were captured by active search and 96 (39.02%) were captured by traps (Table 1).

Different authors report an expressive richness of social wasps in areas of Riparian Forest when compared to other ecosystems (DINIZ & KITAYAMA 1994; CLEMENTE 2009; HENRIQUE-SIMÕES *et al.* 2011; PEREIRA & ANTONIALLI-JUNIOR 2011; LOCHER 2012; BRUNISMANN *et al.* 2016). The Riparian Forests have a large vegetal complexity which may favor social wasps, as it provides a variety of physical supports to nest foundation, rises the amount and heterogeneity of feeding resources, impose less microclimate variability and rises the density of colonies (LAWTON 1983; DINIZ & KITAYAMA 1994; SOUZA *et al.* 2010). Furthermore, these ecosystems are close to hydric resources which may have affected the passive thermal regulation of the colonies. Yet, it is pointed out that the real species richness of this environment might be even higher, as both collection methods resulted in a relatively low proportion of estimated richness when compared to the Restored Forest and Regenerating Cerrado (Table 2).

Regarding social wasps collected in bait traps, 413 specimens (94.7%) belong to Epiponini, 16 (3.7%) to Mischocyttarini and seven (1.6%) to Polistini. In the active search, 328 specimens (68.76%) belong to Epiponini, 133 (27.88%) to Mischocyttarini and 16 (3.35%) to Polistini.

Thus, bait traps were more efficient for collecting wasps of the Epiponini. This result is contrasting to other ones reporting the active search as the most efficient method for Epiponini

in comparison to bait traps (SOUZA & PREZOTO 2006; KLEIN *et al.* 2015). However, these works also resulted in the active search being the most effective to sample species richness of other tribes, as resulted in the present work.

Out of the 163 specimens of *Mischocyttarus monte* Zikán collected at the Restored Forest, 50 (92.6%) were obtained by active search and four (7.4%) by bait traps. A low-capture rate of *Mischocyttarus* species by bait traps were also reported by other studies, such as in SOUZA & PREZOTO (2006), whom worked in Cerrado Fields and Semideciduous Forests. In conserved Ombrophile Forests, no species of this genus was collected by passive methodology, where the bait used was the same as in the present study (SOUZA *et al.* 2012), probably due to the small number of individuals per colony (GIANNOTTI 1998; TORRES *et al.* 2011) hindering the sampling.

In relation to *M. monte*, BRUNISMANN *et al.* (2016) reported it as the most abundant species and its most efficient collection method was active search, highlighting the difficulty of collecting this genus in trap baits.

Thus, it may be concluded that the use of bait traps as the sole method of sampling for social wasps might underestimate the real richness of the Polistini and Mischocyttarini of a given area.

Concerning the most abundant species by each ecosystem, the one with the highest number of collected specimens was the swarming wasp *Agelaia pallipes* Olivier at the Regenerating Cerrado, with 178 recorded specimens, with 117 (39.39%) by bait traps and 61 (20.53%) by active search.

*Agelaia vicina* de Saussure was the most representative in three other ecosystems: at Cerradão, with 20 (15.26%) sampled specimens by traps and 24 (18.32%) by active search; at the Semideciduous Forest, with 23 (30.26%) specimens being sampled by active search and nine (8.4%) by traps; and at the Riparian Forest, with 74 (30.08%) collected individuals by active search and 38 (15.44%) by traps.

At the Restored Forest, out of the 163 collected social vespids, 54 (33.12%) were specimens of *M. monte*, with 50 collected by active search and four by traps. The second-most numerous species was the swarming wasp *Polybia chrysothorax* Lichtenstein, with 41 individuals captured, out of which 39 were captured in bait traps.

In different studies, *A. vicina* was also the most abundant species (SOUZA & PREZOTO 2006; RIBEIRO-JUNIOR 2008; SILVA 2012), possibly due to being the species which builds the largest colonies of Polistinae, as reported by ZUCCHI *et al.* (1995), whom recorded a colony with more than a million specimens of adults and nests with 7.5 million breeding cells.

OLIVEIRA *et al.* (2010) add that a high-growth ratio, a large population and an elevated number of queens is what

**Table 2.** Estimative indexes of richness from Jackknife 1 and 2 using methods of Active Search (AS) and Bait Traps (BT) in five areas from the center-east of the São Paulo State. (RC – Regenerating Cerrado, C – Cerradão, RTF – Restored Forest, SDF – Semideciduous Forest and RPF – Riparian Forest).

Richness	RC		C		RTF		SDF		RPF	
	AS	BT	AS	BT	AS	BT	AS	BT	AS	BT
	13	17	12	9	13	9	10	8	21	14
Estimator J1	18	22	18.66	12.33	16.33	9.4	15.83	13	32.66	21.71
Estimator J2	20.4	24.4	23.46	13.93	17.93	13.4	19.96	17	41.46	26.95
% estimated J1	72.99	77.27	64.30	72.99	79.60	95.74	63.17	61.53	64.29	64.48
% estimated J2	63.72	69.67	51.15	64.60	72.50	67.16	50.10	47.05	50.65	51.94

make nests with such greatness possible to exist. This factor certainly influenced the elevated occurrence of specimens of this group in the active collections and in traps, highlighting the importance of this tribe to the studied ecosystems.

All *Agelaia* species sampled were present in all collection sites. OLIVEIRA et al. (2010) suggest that *A. vicina* should be considered in several environments as a key species, or in other words, as defined by PAINE (1969), a species in which its population determines the stability (integrity and unaltered persistency over time) of a community through its activities and abundances.

Other *Agelaia* species are frequently among the most abundant, be in more opened Cerrados, transition cerrado-forest, forest borders (FERREIRA et al. 2020) or inside forests (KLIEN et al. 2015).

At the sampled phytophysionomies, with exception of Regenerating Cerrado, the active search with entomological net resulted as the most efficient to sample species richness of social wasps. In fact, bait traps collected almost exclusively wasps tribe Epiponini, except for 23 individuals (5.3%) belonging to other tribes.

In relation to abundance of collected individuals, there was a higher efficiency rate of the active search in the Restored Forest, Semideciduous Forest and Riparian Forest. At the Cerradão, both methods resulted as equally efficient, whilst in Regenerating Cerrado the traps resulted as the most efficient method in this context.

The fact that neither of both methods could sample the total of species sampled per each area demonstrated that these strategies are complementary in any sampling and analyses over the social wasp community in different ecosystems. Thus, associating other methodologies, such as Malaise trap (cf. FERREIRA et al. 2020), might contribute to the rise of the captured species number and for a sampling closer to the real richness of these areas.

## ACKNOWLEDGEMENTS

This work received financial support from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). We would like to thank the Instituto de Ecologia (INECOL) A.C. Mexico. for the partnership.

## REFERENCES

Almeida, LM, CS Ribeiro-Costa & L Marioni, 1998. Manual de Coleta, Conservação, Montagem e Identificação de Insetos. Ribeirão Preto, Holos.

Alvarenga, RD, MM Castro, HH Santos-Prezoto & F Prezoto, 2010. Nesting of social wasps (Hymenoptera, Vespidae) in urban gardens in Southeastern Brazil. *Sociobiology*, 55: 445-452.

Aud, AM, CA Carvalho, MA Clemente & F Prezoto, 2010. Diversity of social wasps (Hymenoptera) in a silvipastoral system. *Sociobiology*, 55: 627-636.

Ayres, M, M Ayres Junior, DL Ayres & AA Santos, 2007. BioEstat - Aplicações estatísticas nas áreas das ciências biomédicas. Belém, ONG Mamirauá.

Barbosa, BC, MF Paschoalini & F Prezoto, 2014. Temporal activity patterns and foraging behavior by social wasps (Hymenoptera, Polistinae) on fruits of *Mangifera indica* L. (Anacardiaceae). *Sociobiology*, 61(2): 239-242. DOI: <https://doi.org/10.13102/sociobiology.v61i2.239-242>

Brito, ELS, CA Sá & GMM Santos, 2020. Body size and its relation to the foraging schedules of social wasps. *Neotropical Entomology*, 49: 668-676. DOI: <https://doi.org/10.1007/s13744-020-00789-4>

Brunismann, AG, MM Souza, EP Pires, EL Coelho & LR Milani, 2016. Social wasps (Hymenoptera: Vespidae) in deciduous seasonal forest in Southeastern Brazil. *Journal of Entomology and Zoology Studies*, 4: 447-452.

Bueno, ET, MM Souza & MA Clemente, 2019. The Effect of Forest Fragmentation on Polistinae. *Sociobiology*, 66: 508-514. DOI: <https://doi.org/10.13102/sociobiology.v66i3.4378>

Cardoso-Leite, E, TB Covre, RG Ometto, DC Cavalcanti & MI Pagani, 2004. Fitossociologia e caracterização sucessionária de um fragmento de Mata Ciliar, em Rio Claro, SP, como subsídio à recuperação da área. *Revista do Instituto Florestal*, 16: 31-41.

Carpenter, JM & OM Marques, 2001. Contribuição ao estudo dos vespídeos do Brasil (Insecta, Hymenoptera, Vespidae), Volume 2. Cruz das Almas, Universidade Federal da Bahia, Serie Publicações Digitais.

Clemente, MA, 2009. Vespas Sociais (Hymenoptera, Vespidae) do Parque Estadual do Ibitipoca - MG: Estrutura, Composição e Visitação Floral. Dissertation (Master in Biological Sciences). Universidade Federal de Juiz de Fora.

Clemente, MA, D Lange, W Dáttilo, K Del-Claro & F Prezoto, 2013. Social wasp-flower visiting guild in less structurally complex habitats are more susceptible to local extinction. *Sociobiology*, 60: 337-344. DOI: <https://doi.org/10.13102/sociobiology.v60i3.337-344>

Cooper, MA, 1997. The subgenus *Megacanthopus* Ducke of *Mischocyttarus* de Saussure (Hym., Vespidae) with a key and three new species. *Entomologist's Monthly Magazine*, 133: 217-233.

De Souza, AR, DFA Venâncio, JC Zanuncio & F Prezoto, 2011. Sampling methods for assessing social wasps species diversity in a eucalyptus plantation. *Journal of Economic Entomology*, 104: 1120-1123. DOI: <https://doi.org/10.1603/EC11060>

Diniz, IR & K Kitayama, 1994. Colony densities and preferences for nest habitats of some social wasps in Mato Grosso State, Brazil (Hymenoptera: Vespidae). *Journal of Hymenoptera Research*, 3: 133-143.

Diniz, IR & K Kitayama, 1998. Seasonality of vespidae species (Hymenoptera: Vespidae) in a central Brazilian Cerrado. *Revista de Biologia Tropical*, 46: 109-114.

Elpino-Campos, A, K Del-Claro & F Prezoto, 2007. Diversity of Social Wasps (Hymenoptera, Vespidae) in the Cerrado Fragments of Uberlândia, Minas Gerais State, Brazil. *Neotropical Entomology*, 36: 685-692. DOI: <https://doi.org/10.1590/S1519-566X2007000500008>

Ferreira, JVA, D Storck-Tonon, RJ da Silva, A Somavilla, MJB Pereira & DJ da Silva, 2020. Effect of habitat amount and complexity on social wasps (Vespidae: Polistinae): implications for biological control. *Journal of Insect Conservation*, 2020: 1-12. DOI: <https://doi.org/10.1007/s10841-020-00221-7>

Fragoso, EN, 2005. Ictiofauna da microbacia do córrego da Lapa, bacia do Alto Paraná, Itirapina/Ipeúna, SP. 2005. Thesis (PhD in Ecology and Natural Resources). Universidade Federal de São Carlos.

Giannotti, E, 1998. The colony cycle of the social wasp, *Mischocyttarus cerberus styx* Richards, 1940 (Hymenoptera, Vespidae). *Revista Brasileira de Entomologia*, 41: 217-224.

Hammer, Ø, DAT Harper & PD Ryan, 2001. PAST: Paleontological Statistics software package for education and data analysis. *Palaeontologia Electronica*, 4.

Henrique-Simões, M, MD Cuzzo & FA Friero-Costa, 2011. Social wasps of Unilavras/Boqueirão Biological Reserve, Ingaí, state of Minas Gerais, Brazil. *Check List*, 7: 656-667. DOI: <https://doi.org/10.15560/7.5.656>

Hermes, MG & A Köhler, 2006. The flower-visiting social wasps (Hymenoptera, Vespidae, Polistinae) in two areas of Rio Grande do Sul State, Southern Brazil. *Revista Brasileira de Entomologia*, 50: 268-274. DOI: <https://doi.org/10.1590/>

- S0085-56262006000200008
- Jacques, GC, AA Castro, GK Souza, R Silva-Filho, MM Souza & JC Zanuncio, 2012. Diversity of Social Wasps in the Campus of the “Universidade Federal de Viçosa” in Viçosa, Minas Gerais State, Brazil. *Sociobiology*, 59: 1053-1062. DOI: <https://doi.org/10.13102/sociobiology.v59i3.566>
- Klein, RP, A Somavilla, A Köhler, CV Cademartori & ED Forneck, 2015. Space-time variation in the composition, richness and abundance of social wasps (Hymenoptera: Vespidae: Polistinae) in a forest-agriculture mosaic in Rio Grande do Sul, Brazil. *Acta Scientiarum*, 37: 327-335. DOI: <https://doi.org/10.4025/actasciobiolsci.v37i3.27853>
- Landolt, PJ, CS Smithhisler, HC Reed & LM McDonough, 2000. Trapping social wasps (Hymenoptera: Vespidae) with acetic acid and saturated short chain alcohols. *Journal of Economic Entomology*, 93: 1613-1618. DOI: <https://doi.org/10.1603/0022-0493-93.6.1613>
- Lawton, JH, 1983. Plant architecture and the diversity of phytophagous insects. *Annual Review of Entomology*, 28: 23-39. DOI: <https://doi.org/10.1146/annurev.en.28.010183.000323>
- Lima, ACO, 2008. Sobre a diversidade de vespas sociais (Vespidae: Polistinae) em fragmentos florestais remanescentes do noroeste e do nordeste do Estado de São Paulo, e o seu possível uso como indicadores de conservação da biodiversidade. 2008. Dissertation (Master in Biological Sciences). Universidade de São Paulo.
- Lima, ACO, MSM Castilho-Noll, B. Gomes & FB Noll, 2010. Social wasp diversity (Vespidae, Polistinae) in a forest fragment in the northeast of São Paulo state sampled with different methodologies. *Sociobiology*, 55: 613-623.
- Lima, MAP, JR Lima & F Prezoto, 2000. Levantamento dos gêneros, flutuação das colônias e hábitos de nidificação de vespas sociais (Hymenoptera, Vespidae) no Campus da UFJF, Juiz de Fora, MG. *Revista Brasileira de Zootecias*, 2: 69-80.
- Liow, LH, NS Sodhi & T Elmquist, 2001. Bee diversity along a disturbance gradient in tropical lowland forests of south-east Asia. *Journal of Applied Ecology*, 38: 180-192. DOI: <https://doi.org/10.1046/j.1365-2664.2001.00582.x>
- Locher, GA, 2012. Estudo comparativo da diversidade de vespas sociais (Hymenoptera, Vespidae) em Mata Ciliar e cultura de cana-de-açúcar na região de Ipeúna, SP. Dissertation (Master in Biological Sciences: Zoology). Universidade Estadual Paulista “Júlio de Mesquita Filho”.
- Locher, GA, OC Togni, OT Silveira & E Giannotti, 2014. The social wasp fauna of a riparian forest in southeastern Brazil (Hymenoptera, Vespidae). *Sociobiology*, 61: 225-233. DOI: <https://doi.org/10.13102/sociobiology.v61i2.225-233>
- Marques, OM, CAL Carvalho & JMDa Costa, 1993. Levantamento das espécies de vespas sociais (Hymenoptera: Vespidae) no município de Cruz das Almas – Estado da Bahia. *Insecta*, 2: 1-9.
- Marques, OM, PA Santos, AF Vinhas, ALV Souza, CAL Carvalho & JL Meira, 2005. Social wasps (Hymenoptera: Vespidae) visitors of nectaries of *Vigna unguiculata* (L.) Walp. in the region of Recôncavo of Bahia. *Magistra*, 17: 64-68.
- Melo, AC, GMM Santos, JD Cruz & OM Marques, 2005. Vespas Sociais (Vespidae), pp. 244-257. In: Juncá, FA, L Funch & W Rocha (Eds.). *Biodiversidade e conservação da Chapada Diamantina*. Brasília, Ministério do Meio Ambiente.
- Morato, EF, ST Amarante & OT Silveira, 2008. Avaliação ecológica rápida da fauna de vespas (Hymenoptera, Aculeata) do Parque Nacional da Serra do Divisor, Acre, Brasil. *Acta Amazonica*, 38: 789-798. DOI: <https://doi.org/10.1590/S0044-59672008000400025>
- Neto, SS, RC Monteiro, RA Zucchi & RCB Moraes, 1995. Uso da análise faunística de insetos na avaliação do impacto ambiental. *Scientia Agricola*, 52: 9-15. DOI: <https://doi.org/10.1590/S0103-90161995000100003>
- Oliveira, SAD, MM Castro & F Prezoto, 2010. Foundation pattern, productivity and colony success of the paper wasp, *Polistes versicolor*. *Journal of Insect Science*, 10: 125. DOI: <https://doi.org/10.1673/031.010.12501>
- Pagano, SN & HF Leitão-Filho, 1987. Composição florística do estrato arbóreo de mata mesófila semidecídua no município de Rio Claro (estado de São Paulo). *Revista Brasileira de Botânica*, 10: 3-47.
- Pagano, SN, HF Leitão-Filho & GJ Shepherd, 1987. Estudo fitossociológico em mata mesófila semidecídua no município de Rio Claro (estado de São Paulo). *Revista Brasileira de Botânica*, 10: 49-62.
- Paine, RT, 1969. A note on trophic complexity and community stability. *The American Naturalist*, 103: 91-93. DOI: <https://doi.org/10.1086/282586>
- Pereira, MGC & WF Antoniali-Junior, 2011. Social wasps in riparian forest in Batayporã, Mato Grosso do Sul State, Brazil. *Sociobiology*, 57: 153-163.
- Pickett, KM & JW Wenzel, 2007. Revision and cladistic analysis of the nocturnal social wasp genus, *Apoica* Lepeletier (Hymenoptera: Vespidae; Polistinae, Epiponini). *American Museum Novitates*, 2007: 1-30. DOI: [https://doi.org/10.1206/0003-0082\(2007\)397\[1:RACAOT\]2.0.CO;2](https://doi.org/10.1206/0003-0082(2007)397[1:RACAOT]2.0.CO;2)
- R Development Core Team, 2009. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Available in: <http://www.R-project.org>.
- Raveret-Richter, M, 2000. Social wasp (Hymenoptera: Vespidae) foraging behavior. *Annual Review of Entomology*, 45: 121-150. DOI: <https://doi.org/10.1146/annurev.ento.45.1.121>
- Ribeiro, JF & BMT Walter, 1998. As principais fitofisionomias do Bioma Cerrado, pp. 152-212 In: Sano, SM, SP Almeida & JF Ribeiro (Eds.). *Ecologia e flora – vol.1*. Brasília, EMBRAPA.
- Ribeiro-Junior, C, 2008. Levantamento de vespas sociais (Hymenoptera: Vespidae) em uma Eucaliptocultura. Dissertation (Master in Biological Sciences). Universidade Federal de Juiz de Fora.
- Richards, OW, 1978. *The social Wasps of the Americas, Excluding the Vespinae*. London, British Museum, Natural History.
- Rizzini, CT, 1997. *Tratado de Fitogeografia do Brasil: aspectos ecológicos, sociológicos e florísticos*. Rio de Janeiro, Âmbito Cultural Edições Ltda.
- Rodrigues, VM & VLL Machado, 1982. Vespídeos sociais: Espécies do Horto Florestal “Navarro de Andrade” de Rio Claro, SP. *Naturalia*, 7: 173-175.
- Santos, BB, 1996. Ocorrência de vespídeos sociais (Hymenoptera, Vespidae) em pomar em Goiânia, Goiás, Brasil. *Agrárias*, 15: 43-46.
- Santos, GMM, FCC Bichara, JJ Resende, JD Cruz & OM Marques, 2007. Diversity and community structure of social wasps (Hymenoptera: Vespidae) in three ecosystems in Itaparica island, Bahia State, Brazil. *Neotropical Entomology*, 36: 180-185. DOI: <https://doi.org/10.1590/S1519-566X2007000200002>
- Santos, GMM, JD Cruz, OM Marques & N Gobbi, 2009. Diversidade de vespas sociais (Hymenoptera: Vespidae) em áreas de cerrado na Bahia. *Neotropical Entomology*, 38: 317-320. DOI: <https://doi.org/10.1590/S1519-566X2009000300003>
- Santos, GMM & SJ Presley, 2010. Niche Overlap and Temporal Activity Patterns of Social Wasps (Hymenoptera: Vespidae) in a Brazilian Cashew Orchard. *Sociobiology*, 56: 121-131.
- Silva, NJ, 2012. Diversidade de vespas sociais em cultivo de cana-de-açúcar. Dissertation (Master in Ecology). Universidade Federal de Juiz de Fora.
- Silveira, OT, 2002. Surveying Neotropical social wasps. An evaluation of methods in the “Ferreira Penna” Research station (ECFPn), in Caxiuanã, PA, Brazil (Hym., Vespidae, Polistinae). *Papéis Avulsos de Zoologia*, 42: 299-323. DOI: <https://doi.org/10.1590/S0031-10492002001200001>
- Silveira, OT, MC Esposito, JN Santos Júnior & FE Gemaque

- Júnior, 2005. Social wasps and bees captured in carrion traps in a rain forest in Brazil (Hymenoptera: Vespidae; Apidae). *Entomological Science*, 8: 33-39. DOI: <https://doi.org/10.1111/j.1479-8298.2005.00098.x>
- Siqueira, LP, 2002. Monitoramento de áreas restauradas no interior do Estado de São Paulo, Brasil. Dissertation (Master in Forest Resources). Universidade de São Paulo.
- Somavilla, A, DWA Marques, EAS Barbosa, JSP Junior & ML Oliveira, 2014. Vespas Sociais (Vespidae: Polistinae) em uma Área de Floresta Ombrófila Densa Amazônica no Estado do Maranhão, Brasil. *EntomoBrasilis*, 7: 183-187. DOI: <https://doi.org/10.12741/ebrasilis.v7i3.404>
- Somavilla, A, RNM Moraes Junior & JA Rafael, 2019. Is the social wasp fauna in the tree canopy different from the understory? Study of a particular area in the Brazilian Amazon Rainforest. *Sociobiology*, 66: 79-185. DOI: <https://doi.org/10.13102/sociobiology.v66i1.3568>
- Souza, MM & F Prezoto, 2006. Diversity of social wasps (Hymenoptera: Vespidae) in semideciduous forest and cerrado (Savanna) regions in Brazil. *Sociobiology*, 47: 135-147.
- Souza, MM, TE Ladeira, NRG Assis, A Elpino-Campos, P Carvalho & JNC Louzada, 2010. Ecologia de vespas sociais (Hymenoptera, Vespidae) no Campo Rupestre na Área de Proteção Ambiental, APA, São José, Tiradentes, MG. *MG.Biota*, 3: 15-30.
- Souza, MM, EP Pires, M Ferreira, TE Ladeira, M Pereira, A Elpino-Campos & JC Zanuncio, 2012. Biodiversidade de vespas sociais (Hymenoptera: Vespidae) do Parque Estadual do Rio Doce, Minas Gerais, Brasil. *MG.Biota*, 5: 4-19.
- Souza, MM, LN Perillo, BC Barbosa & F Prezoto, 2015. Use of flight interception traps of Malaise type and attractive traps for social wasps record (Vespidae: Polistinae). *Sociobiology*, 62: 450-456. DOI: <https://doi.org/10.13102/sociobiology.v62i3.708>
- Souza, MM, GS Teofilo-Guedes, LR Milani, ASB Souza & PP Gomes, 2020. Social Wasps (Vespidae: Polistinae) from the Brazilian Atlantic Forest. *Sociobiology*, 67: 1-12. DOI: <https://doi.org/10.13102/sociobiology.v67i1.4597>
- Tanaka-Junior, GM & FB Noll, 2011. Diversity of Social Wasps on Semideciduous Seasonal Forest Fragments with Different Surrounding Matrix in Brazil. *Psyche*, 2011: 861747. DOI: <https://doi.org/10.1155/2011/861747>
- Togni, OG, 2009. Diversidade de vespas sociais (Hymenoptera, Vespidae) na Mata Atlântica do litoral norte do Estado de São Paulo. Dissertation (Master in Biological Sciences: Zoology). Universidade Estadual Paulista "Júlio de Mesquita Filho".
- Togni, OC, GA Locher, E Giannotti & OT Silveira, 2014. The social wasp community (Hymenoptera, Vespidae) in an area of Atlantic Forest, Ubatuba, Brazil. *Check List*, 10: 10-17. DOI: <https://doi.org/10.15560/10.1.10>
- Torres, VO, TS Montagna, WD Fernandes & WF Antonialli-Junior, 2011. Colony cycle of the social wasp *Mischocyttarus consimilis* Zikán (Hymenoptera, Vespidae). *Revista Brasileira de Entomologia*, 55: 247-252. DOI: <https://doi.org/10.1590/S0085-56262011000200016>
- Wegner, GS & KK Jordan, 2005. Comparison of three liquid lures for trapping social wasps (Hymenoptera: Vespidae). *Journal of Economic Entomology*, 98: 664-666. DOI: <https://doi.org/10.1603/0022-0493-98.3.664>
- Zucchi, R, SF Sakagami, FB Noll, MR Mechi, S Mateus, MV Baio & SN Shima, 1995. *Agelais vicina*, a swarm-founding Polistinae with the largest colony size among wasps and bees (Hymenoptera: Vespidae). *Journal of the New York Entomological Society*, 103: 129-137.

\*\*\*\*\*

